


Results 2014



Legend to indicators

1. Proportion of men who were reported to NPCR within 30 days of their cancer diagnosis.
2. Proportion of men who have a named clinical nurse specialist.
3. Proportion of men who had a first outpatient visit due to suspicion of prostate cancer within 60 days of referral.
4. Proportion of men who were informed of their cancer diagnosis within 18 days after prostate biopsy.
5. Proportion of men up to 80 years of age with high risk cancer who were investigated for bone metastases. High risk cancer: T1,2 and Gleason 8-10 or PSA 20-50 ng/ml or T3 and PSA < 50 ng/ml.
6. Proportion of men with very low risk cancer who were started on active surveillance. Cancer with very low risk: T1c, Gleason 6 or lower, PSA below 10 ng/ml, PSA density below 0.15 ng/ml, no more than four cores with cancer, no more than 8 mm of cancer in total at biopsy.
7. Proportion of men with high risk cancer for whom curative Tx can be considered who were discussed at a multidisciplinary team meeting.
8. Proportion of men not older than 75 years with localized high risk cancer who received curative treatment. Localized high risk cancer: T1,2 and Gleason 8-10 and/or PSA 20-50 ng/ml and no metastases.
9. Proportion of men for whom nerve sparing intention at radical prostatectomy was documented preoperatively .
10. Proportion of men diagnosed with pT2 cancer at prostatectomy who had negative margins, i.e. radically removed tumor.

Symbols

Quality indicators in NPCR and three levels of performance

- 2** High level
- 1** Intermediate level
- 0** Low level

Patients

Number of patients for whom high quality level was reached out of all reported patients.

My clinic/Sweden overall

Number of patients for whom the indicator level was reached. Vertical lines indicate upper and lower cut-offs. Mean value for the whole country.

Performance

Performance in the last four years.

Would you like to know more? Go to www.npcr.se where the annual report is available